

Article

Tocilizumab and sarilumab

Tocilizumab was approved for use in patients with RA in 2009, making it a more recent biologic drug, with sarilumab being the most recent, having been approved in 2017.

Print

Original Biologic drug Method of administration

Tocilizumab intravenous infusion, once every 4 weeks or weekly

subcutaneous (under the skin) injection

Sarilumab subcutaneous (under the skin) injection every other week

How does it work?

As with other biologic drugs, tocilizumab and sarilumab work by targeting proteins called cytokines, which are responsible for the inflammation caused by the immune system's response. In this case the cytokines being targeted are called 'IL6 receptors'.

Most commonly reported side effects

As with any medication, tocilizumab has a number of possible side effects, although it is important to remember that these are only potential side effects. They may not occur at all.

Common side effects may include:

Tocilizumab and sarilumab:

- upper respiratory tract infections with typical symptoms such as cough, blocked nose, runny nose, sore throat and headache
- high blood fat (cholesterol) levels
- neutropenia (low neutrophils, which are a type of white blood cell)
- increased liver enzymes (an indication that the liver is affected)
- injection site reactions
- upper respiratory infections and urinary tract infections

More information on side effects can be found in the patient information leaflet for tocilizumab. Remember to report any concerns about possible side effects to the doctors and nurses.

IL6 inhibitors with other medicines

Some biologic drugs are known to interact poorly with other biologics. You may therefore be asked to leave a gap between stopping one biologic drug and starting another, so that the first drug has time to start coming out of your system.

Tocilizumab is known to interact poorly with the anti-psychotic drug 'clozapine'.

IL6 inhibitors during pregnancy and breastfeeding

Tocilizumab and sarilumab should not be used during pregnancy unless clearly necessary,

Tocilizumab and sarilumab should not be used during pregnancy or while breastfeeding, due to limited data on safety. There is also not enough data to suggest that men can take this drug while their partner is trying to conceive.

Pregnancy information in this booklet is based on British Society for Rheumatology (BSR) guidelines on prescribing drugs in pregnancy and breastfeeding.

Before starting a family it is recommended that you get advice from the consultant or clinical nurse specialist about when to start a pregnancy.

IL6 inhibitors and alcohol

You can drink alcohol on these medications. However, it is not uncommon when taking a biologic drug to be on other medications, where different guidance applies. Methotrexate, for example, can affect the liver, so for those taking methotrexate alongside their biologic, moderate intake of alcohol is recommended in line with government guidelines.

IL6 inhibitors and immunisations/ vaccinations

Live vaccines (measles, mumps, rubella, i.e. MMR, chickenpox, oral polio (NOT injectable Live vaccines (measles, mumps, rubella i.e. MMR, chickenpox, oral polio (NOT injectable polio), BCG, oral typhoid and yellow fever) cannot be given to anyone already taking tocilizumab or sarilumab. If the treatment has not yet been started, it is important to seek advice on how long a gap to leave after having a live vaccine.

Medicines in rheumatoid arthritis

We believe it is essential that people living with RA understand why certain medicines are used, when they are used and how they work to manage the condition.

Order/Download



Updated: 01/09/2020

This article was downloaded from www.nras.org.uk. National Rheumatoid Arthritis Society (NRAS) is a registered charity in England and Wales (1134859) and Scotland (SC039721). A private company limited by guarantee. Registered in England and Wales (7127101).